

Sample Questions for Chem 002 Final FS10

1. MSDS (the rest listed on review):

- a. Proper attire –
- b. Acid Spill –
- c. Bunsen Burners –
- d. Phenolphthalein –
- e. Types of radiation (listed below) are stopped by what type of material?
 - alpha –
 - beta –
 - gamma –
 - neutron –

2. Ternary Mixture – A mixture is known to contain the four compounds in the table.

- a. Draw a flow chart to show the steps that you would use to separate the following compounds.

	Cold water	Hot water	3M HCl	3M NaOH
benzoic acid	no	yes	no	yes
Mg(OH) ₂	no	no	yes	no
Na ₂ SO ₄	yes	yes	yes	yes
Zn(OH) ₂	no	no	yes	yes

- b. The initial mass was 5.025g. The resulting masses were benzoic acid = 1.760g, Mg(OH)₂ = 0.754g, Na₂SO₄ = 1.005g, and Zn(OH)₂ 1.256g. Calculate the percent recovery of each component and the total percent recovery.

3. Radioactive Decay:

a. Balance the following radioactive decay equations:



b. Determine the specific decay constant, initial activity and half-life of a radioactive isotope. Given the equations:

$$A = A_0 e^{-kt} \quad \ln A = -kt + \ln A_0 \quad \ln 2 = 0.693 \quad t_{1/2} = \ln 2 / k \quad y = mx + b \quad m = (y_2 - y_1) / (x_2 - x_1)$$

and the data:

Time, minutes	Counts/Min	ln (Counts/Min)
0		
2	14472	
3	14328	
4	14248	
5	14095	
6	13920	
10	13359	

1. Determine the specific decay constant, k , for this radioactive decay.

2. Determine the initial activity, A_0 .

3. Determine the half-life.

4. Heat of Neutralization:

A reaction of 100mL of 1.35M HCl and 100mL of 1.76M NaOH is monitored and the following temperatures were recorded: starting temperature = 24.6 °C; and final temperature = 38.8 °C. Calculate the ΔH of this reaction.

Given: C_p of solution (J/K) = 4.13 J/(g · K) * Volume of solution in mL (1 mL \approx 1 g for aqueous soln)

C_p of calorimeter (J/K) = 50

$Q = (-\text{total } C_p * \Delta T)$

$\Delta H = Q/n$

$\Delta T = T_f - T_i$

$n = \#$ of moles reacted

- Determine the change in temperature for the system.
- Determine the C_p of the solution (J/K).
- Determine the **total** C_p of the system.
- Determine the number of moles of the acid and the base. Which is the **limiting reagent**?
- Determine the **Heat Transfer, Q**, for the reaction.
- Determine the **change in enthalpy, ΔH** , for the reaction.

5. Heat of Fusion. An ice cube with mass 9.53 grams (presume $T_i = 0\text{ }^\circ\text{C}$) is placed in a calorimeter containing 111.24 grams of distilled water at a temperature of $23.2\text{ }^\circ\text{C}$. After equilibration, the final temperature was $15.8\text{ }^\circ\text{C}$.

Given: $\Delta H_{\text{total}} = \Delta H_{\text{ice}} + \Delta H_{\text{water}} + \Delta H_{\text{calorimeter}} + \Delta H_{\text{fus}} = 0$

$$\Delta H_{\text{water}} (\text{J}) = C_{p,\text{H}_2\text{O}} * (\text{mass}) * \Delta T$$

$$\Delta H_{\text{ice}} (\text{J}) = C_{p,\text{ice}} * (\text{mass}) * \Delta T$$

$$\Delta H_{\text{calorimeter}} (\text{J}) = C_{p,\text{Cal}} * \Delta T$$

$$C_{p,\text{H}_2\text{O}} = 4.18 \text{ J}/(\text{g}^\circ\text{C})$$

$$C_{p,\text{Cal}} = 50 \text{ J}/^\circ\text{C}$$

$$\Delta T = T_f - T_i$$

a. Determine the ΔH_{water} .

b. Determine the ΔH_{ice} . (*Hint for C_p – The ice has melted.*)

c. Determine the $\Delta H_{\text{calorimeter}}$.

d. Determine the ΔH_{fus} for **one gram** of ice.

(*Hint:* For a calorimeter (i.e., closed systems) $\Delta H_{\text{total}} = 0$)

f. If heat transfers from the system (solute) to the surroundings (solvent), then ΔH is negative ($\Delta H < 0$), and the reaction is defined as (endothermic / exothermic) and the temperature of the solvent will go (up / down).

g. If heat transfers from the surroundings (solvent) to the system (solute), then ΔH is positive ($\Delta H > 0$), and the reaction is defined as (endothermic / exothermic) and the temperature of the solvent will go (up / down).

h. The heat of neutralization experiment was an (endothermic / exothermic) reaction .

i. The heat of fusion experiment was an (endothermic / exothermic) reaction.

j. This term means “the techniques that are used to measure enthalpy”:

k. This term means “the energy needed to raise the temperature of an object 1°C ”:

l. This term means “the energy needed to raise the temperature of one gram of a substance 1°C ”:

m. The heat capacity is an extrinsic property. Define intrinsic and extrinsic properties and give an example of each.

6. Antacids: You are given 1.12 M HCl and 1.48 M NaOH. The antacid you use contains 300 mg of CaCO_3 and 100 mg of Al(OH)_3 . If the antacid dissolved in 35.0 ml of HCl and was then back titrated with 21.8 ml of NaOH, find the following:

- a. The original number mmoles of HCl used to dissolve the antacid and neutralize the base.

- b. The number of mmoles of NaOH used to backtitrate the acid.

- c. The number of mmoles of **acid*** used to neutralize only the antacid (a.k.a. the excess HCl).
(*Note: Handout dated 110418 had antacid here, should have been acid.)

- d. Write the balanced equations for the neutralization of the antacid (Both CaCO_3 and Al(OH)_3).

- e. Using the number of mg in the tablet, calculate the mmoles of each component
(Both CaCO_3 and Al(OH)_3).

- f. Based on the mmoles of each component, calculate the theoretical number of mmoles of HCl that should have been needed to neutralize the antacid. (*Hint: Use the mole ratios.*)

- g. What was the total number of theoretical mmoles of HCl that should have been neutralized?

- h. Compare the theoretical (g.) to the actual (c.). What are possible reasons this discrepancy could have occurred?

7. Spectrophotometry: Using a Spectrophotometer (Spec 20), a student recorded below the Percent Transmittance data for the following solutions:

**Red Dye Standard (6.30 ppm)
Blue Dye Standard (5.05 ppm)
Purple Unknown**

	400 nm	450 nm	500 nm	550 nm	600 nm	650 nm
Red Std	63.5	48.5	23.5	38.6	78.3	98.5
Blue Std	80.5	99.0	82.5	56.5	8.5	72.4
Purple Unk	79.3	72.5	35.5	85.8	45.5	65.3

a. Calculate the Absorbance for each of the %T listed above .

	400 nm	450 nm	500 nm	550 nm	600 nm	650 nm
Red Std						
Blue Std						
Purple Unk						

b. Determine the following from the data calculated in Part 1 (2 pts):

Red Dye Max. Absorbance = _____ at _____ nm (λ Max)


Blue Dye Max. Absorbance = _____ at _____ nm (λ Max)

c. Calculate the Absorbance Ratio of the Unknown/Standard at (λ Max).

d. Calculate the Dye Concentration in the Unknown. (Standard Concentrations given above.)

	Abs of Unknown (at λ Max)	Abs of Standard (at λ Max)	Abs Ratio Unk/Std (at λ Max)	Dye Conc. in Unknown
Red in Purple				
Blue in Purple				

8. Colorimetry: Using the well strips below, the student put the following number of drops in the wells. In strips I & II, 1-8 drops of blue dye standard solution (5.05 ppm) were added as shown in the diagram. In strip II, additional drops of water were added in order to have the same total volume of 8 drops for each well.



Given: The student found that the unknown solution of blue dye matched well #7 on Strip II.

- What is changing in the first well strip – concentration or pathlength?
- What is changing in the second well strip – concentration or pathlength?
- Looking from the top how does the intensity compare for Strip 1 to Strip 2?
more intense – the same – less intense
- Using $C_1V_1 = C_2V_2$, what is the approximate concentration in ppm for the unknown?

9. Gas Laws: Using the ideal gas law calculate the volume of the system.

Given: pressure = 738 mmHg, mass = 0.725 grams, $MW_{\text{butane}} = 58.000 \text{ g/mole}$, $T = 20^\circ\text{C}$, $R = 0.08206 \text{ Latm/molK}$

a. What is the number of moles of butane?

b. What is the pressure in atm?

c. What is the temperature in K?

d. What is the volume of the system?

e. What would the volume be at STP?

10. Statistics:

a. For the following data set (2.10, 3.20, 3.50, 4.90, 4.30, 2.90) find the mean (average).

b. For the average of the data set above, calculate the % Error if the expected answer was 3.500.

b. For this data set would you calculate the standard deviation or the standard deviation estimate? Explain why.

11. Dimensional Analysis: Choose problems from sets 1, 2, 4 or 5 and work them.

12. Scientific Notation & Significant Figures:

a. Choose problems from sets 1 & 2 and work them.

b. Review problems from the midterm exam.

****Note:** Most of the questions on the final will be similar to those on review and on quizzes.

13. People – How did these people contribute to the experiments we did in Chem 2?

- a. Henri Becquerel (Nuclear)
- b. Pierre and Marie Curie (Nuclear)
- c. Ernst Rutherford (Nuclear)
- d. Albert Einstein (*Nuclear*)
- e. Svante Arrhenius (Antacid)
- f. Johannes Nicolaus Brønsted and Thomas Martin Lowry (Antacid)
- g. Gilbert N. Lewis (Antacid)
- h. Robert Boyle (Gas Laws)
- i. Jacques-Alexandre Charles (Gas Laws)
- j. Amedeo Avogadro (Gas Laws)
- k. Joseph-Louis Gay-Lussac (Gas Laws)
- l. John Dalton (Gas Laws)
- m. Johannes Diderik van der Waals (Gas Laws)
- n. My TA's name is...

14. Glassware and equipment: Identify the equipment below.

